

On K_p -series and varieties generated by wreath products of p -groups

Vahagn H. Mikaelian

To Professor Boris I. Plotkin on his 90'th anniversary

ABSTRACT. Let A be a nilpotent p -group of finite exponent, and B be an abelian p -groups of finite exponent. Then the wreath product $A \operatorname{Wr} B$ generates the variety $\operatorname{var}(A) \operatorname{var}(B)$ if and only if the group B contains a subgroup isomorphic to the direct product $C_{p^v}^\infty$ of countably many copies of the cycle C_{p^v} of order $p^v = \exp(B)$. The obtained theorem continues our previous study of cases when $\operatorname{var}(A \operatorname{Wr} B) = \operatorname{var}(A) \operatorname{var}(B)$ holds for some other classes of groups A and B (abelian groups, finite groups, etc.).

1. Introduction and background information

Nilpotent and locally nilpotent subgroups are proven to be very efficient means to study groups. In particular, the Fitting subgroup and the Frattini subgroup help in study of finite groups, while the Plotkin-Hirsch radical allows to consider infinite groups (see [5, 6, 26, 27, 28, 8, 29] and literature cited therein). This trend is naturally inherited by varieties of groups: nilpotent varieties are in some sense simpler varieties (they are irreducible, have finite base rank, their finitely generated groups are factors of finite powers of relatively free groups, etc.), and investigation of nilpotent subvarieties of a variety is an approach to study general varieties. In the current note we use sequences of nilpotent subvarieties to study varieties generated by non-nilpotent wreath products of groups (here we assume Cartesian wreath products, although the analogs of the statements below also are true for direct wreath products).

Let us introduce the general context in which we examine varieties generated by wreath products. One of the most efficient methods to study product varieties $\mathfrak{U}\mathfrak{V}$ is finding some groups A and B such that $\mathfrak{U} = \operatorname{var}(A)$, $\mathfrak{V} = \operatorname{var}(B)$, and the wreath product $A \operatorname{Wr} B$ generates $\mathfrak{U}\mathfrak{V}$, that is, when the equality

$$(*) \quad \operatorname{var}(A \operatorname{Wr} B) = \operatorname{var}(A) \operatorname{var}(B)$$

holds for the given A and B . Indeed, the product $\mathfrak{U}\mathfrak{V}$ consists of all possible extensions of all groups $A \in \mathfrak{U}$ by all groups $B \in \mathfrak{V}$. If $(*)$ holds for some fixed groups A and B , generating the varieties $\mathfrak{U}$ and $\mathfrak{V}$, then one can restrict to consideration of $\operatorname{var}(A \operatorname{Wr} B)$, which is easier to study rather than to explore all the extensions inside $\mathfrak{U}\mathfrak{V}$. Examples of application of this approach are numerous: for earliest results and references see Chapter 2 of Hanna Neumann's monograph [23] and the work of

1991 *Mathematics Subject Classification*. Primary: 20E22. Secondary: 20E10, 20K01, 20K25, 20D15.

Key words and phrases. Wreath products, varieties of groups, products of varieties of groups, nilpotent varieties, abelian groups, nilpotent groups, K_p -series, nilpotent wreath products, p -groups.

The author was supported in part by joint grant 15RF-054 of RFBR and SCS MES RA, and by 15T-1A258 grant of SCS MES RA.

This article reflects the main plenary talk at the *Groups, Algebras and Identities* conference honoring Professor Boris I. Plotkin's 90th birthday in Jerusalem, Israel, March 20-24, 2016.

G. Baumslag, R.G. Burns, G. Higman, C. Haughton, B.H. Neumann, H. Neumann, P.M. Neumann [4, 2, 7], etc.

This motivated our systematic study of equality (*) for as wide classes of groups as possible. In [16]–[18] we gave a complete classification of all cases, when (*) holds for *abelian* groups A and B , and in [19] and [20] we fully classified the cases when A and B are any *finite* groups.

In the current note we consider the case, when A and B are any p -groups of finite exponents, A is nilpotent and B is abelian. This is a part of larger, only partially published research on cases when (*) holds for nilpotent and abelian groups with some conditions (see also [21]). The main theorem of this note is:

THEOREM 1. *Let A be a non-trivial nilpotent p -group of finite exponent, and B be a non-trivial abelian group of finite exponent p^v . Then the wreath product $A \operatorname{Wr} B$ generates the variety $\operatorname{var}(A) \operatorname{var}(B)$, that is, the variety $\operatorname{var}(A) \mathfrak{A}_{p^v}$ if and only if the group B contains a subgroup isomorphic to the direct product $C_{p^v}^\infty$ of countably many copies of the cyclic group C_{p^v} of order p^v .*

Since B is a non-trivial group of finite exponent, $v > 0$ and by Prüfer-Kulikov's theorem [30, 10] it is a direct product of copies of some finite cyclic subgroups of prime-power orders. The theorem above states that in this direct product the cycles of order p^v must be present at least *countably many* times, whereas the number of direct summands of orders $p^{v-1}, p^{v-2}, \dots, p$ is of no importance.

We below without any definitions use the basic notions of the theory of varieties of groups such as varieties, relatively free groups, discriminating groups, etc. All the necessary definitions and background information can be found in Hanna Neumann's monograph [23]. Following the conventional notation, we denote by $\mathbf{Q}\mathfrak{X}$, $\mathbf{S}\mathfrak{X}$, $\mathbf{C}\mathfrak{X}$ and $\mathbf{D}\mathfrak{X}$, the classes of all homomorphic images, subgroups, Cartesian products and of direct products of finitely many groups of $\mathfrak{X}$, respectively. By Birkhoff's Theorem [3, 23] for any class of groups $\mathfrak{X}$ the variety $\operatorname{var}(\mathfrak{X})$ generated by it can be obtained from $\mathfrak{X}$ by three operations: $\operatorname{var}(\mathfrak{X}) = \mathbf{QSC}\mathfrak{X}$. For information on wreath products we refer to [23, 10]. For the given classes of groups $\mathfrak{X}$ and $\mathfrak{Y}$ we denote $\mathfrak{X} \operatorname{Wr} \mathfrak{Y} = \{X \operatorname{Wr} Y \mid X \in \mathfrak{X}, Y \in \mathfrak{Y}\}$. The specific notions, related to K_p -series and to nilpotent wreath products can be found in [12, 14, 31, 32, 13] and in Chapter 4 of J.D.P. Meldrum's monograph [15].

When this work was in progress we discussed the topic with Prof. A.Yu. Ol'shanskii, who suggested ideas for an alternative (very different from our approach) proof for Theorem 1 using the arguments of his work [24]. In [21] we present the outlines of both proofs side by side.

2. The K_p -series and the proof for Theorem 1

In spite of the fact that the soluble wreath products are “many” (the wreath product of any soluble group is soluble), the nilpotent wreath products are “fewer”: as it is proved by G. Baumslag in 1959, a Cartesian or direct wreath product of non-trivial groups A and B is nilpotent if and only if A is a nilpotent p -group of finite exponent p , and B is a finite p -group [1].

Even after having such an easy-to-use criterion to detect, if the given wreath product is nilpotent, it turned out to be a much harder task and took almost two decades to explicitly compute its nilpotency class in general case. H. Liebeck started by consideration of the cases of wreath products of abelian groups [12], and the final general

formula was found by D. Shield's work [31, 32] in 1977. Later the proof was much shortened by R. Marconi [13].

In order to write down the formula we need the notion of K_p -series. For the given group G and the prime number p the K_p series $K_{i,p}(G)$ of G is defined for $i = 1, 2, \dots$ by:

$$(1) \quad K_{i,p}(G) = \prod_{r,j \text{ with } rp^j \geq i} \gamma_r(G)^{p^j},$$

where $\gamma_r(G)$ is the r 'th term of the lower central series of G .

In particular, $K_{1,p}(G) = G$ holds for any G . From definition it is clear that a K_p series is a descending series, although it may not be strictly descending: some of its neighbor terms may coincide. If G is abelian, then in (1) the powers $\gamma_1(G)^{p^j} = G^{p^j}$ of the first term $\gamma_1(G) = G$ need be considered only.

EXAMPLE 1. If $G = C_{p^3} \times C_p \times C_p$ with $p = 5$, then it is easy to calculate that:

$$\begin{aligned} K_{1,p}(G) &= G; \quad K_{2,p}(G) = \dots = K_{5,p}(G) \cong C_{p^2}; \\ K_{6,p}(G) &= \dots = K_{25,p}(G) \cong C_p; \quad K_{26,p}(G) \cong \{1\}. \end{aligned}$$

If G is some finite p -group, using the K_p -series one may introduce the following additional parameters: let d be the maximal integer such that $K_{d,p}(G) \neq \{1\}$. Then for each $s = 1, \dots, d$ define $e(s)$ by

$$p^{e(s)} = |K_{s,p}/K_{s+1,p}|,$$

and set a and b by the rules:

$$a = 1 + (p-1) \sum_{s=1}^d (s \cdot e(s)), \quad b = (p-1)d.$$

The above integer d does exist, and our notations are correct, for, a finite p -group is nilpotent, its K_p -series will eventually reach the trivial subgroup. To keep the notations simpler, the initial group G is not included in the notations of d , $e(s)$, a and b . But from the context it will always be clear which group G is being considered.

Let A be a nilpotent p -group of exponent p^u , and B be an abelian group of exponent p^v , with $u, v > 0$. Assume all the parameters d , $e(s)$, a , b are defined specifically for the group B . Then by Shield's formula [32] (see also Theorem 2.4 in [15]) the nilpotency class of the wreath product $A \text{ Wr } B$ is the maximum

$$(2) \quad \max_{h=1, \dots, c} \{a h + (s(h) - 1)b\},$$

where $s(h)$ is defined as follows: $p^{s(h)}$ is the exponent of the h 'th term $\gamma_h(A)$ of the lower central series of A .

EXAMPLE 2. If (again for $p = 5$) B is the group $C_{p^3} \times C_p \times C_p$ mentioned in previous example, and if A is the group, say, $C_{p^2} \times C_p$, then $d = 25$; $e(1) = 3$, $e(2) = e(3) = e(4) = 0$, $e(5) = 1$, $e(6) = \dots = e(24) = 0$, $e(25) = 1$, $e(26) = 0$; $a = 133$; $b = 100$; $c = 1$ and $s(1) = 2$. Thus, the nilpotency class of the wreath product $A \text{ Wr } B = (C_{p^2} \times C_p) \text{ Wr } (C_{p^3} \times C_p \times C_p)$ in this case is equal to

$$a \cdot 1 + (s(1) - 1)b = 133 + (2 - 1)100 = 233.$$

In order to prove Theorem 1 we will apply Shield's formula to two auxiliary groups. To construct the first group denote by β the cardinality of B and by A^β the Cartesian product of β copies of A . For the given fixed positive integer l and for the integer $t \geq l$ introduce the group $Z(l, t)$ as the direct product:

$$(3) \quad Z(l, t) = \underbrace{C_{p^v} \times \cdots \times C_{p^v}}_l \times \underbrace{C_{p^{v-1}} \times \cdots \times C_{p^{v-1}}}_{t-l}.$$

LEMMA 1. Assume A , B and β are defined as above and l is any positive integer. Also, assume the exponent of $\gamma_c(A)$ is p^α ($\alpha \neq 0$, since the class of A is c). Then there is a positive integer t_0 such that for all $t > t_0$ the nilpotency class of the wreath product $A^\beta \text{Wr } Z(l, t)$ is equal to

$$(4) \quad c + ct(p-1) \left(\frac{1-p^{v-1}}{1-p} + l/t \cdot p^{v-1} \right) + (\alpha-1)(p-1)p^{v-1}.$$

PROOF. Denote $Z = Z(l, t)$ and notice that $A^\beta \text{Wr } Z$ is nilpotent by Baumslag's theorem [1]. Let us compute the K_p -series for Z and, to keep the notations simpler, not include in the formulas the underbraces of (3) with l and $t-l$. For $i = 1$ we have $K_{1,p}(Z) = Z$. For $i = 2, \dots, p$ we get:

$$K_{i,p}(Z) = Z^p = C_{p^{v-1}} \times \cdots \times C_{p^{v-1}} \times C_{p^{v-2}} \times \cdots \times C_{p^{v-2}}.$$

For $i = p^k + 1, \dots, p^{k+1}$ we have:

$$K_{i,p}(Z) = Z^{p^{k+1}} = C_{p^{v-(k+1)}} \times \cdots \times C_{p^{v-(k+1)}} \times C_{p^{v-(k+2)}} \times \cdots \times C_{p^{v-(k+2)}}.$$

In particular, for $i = p^{v-3} + 1, \dots, p^{v-2}$ we get:

$$K_{i,p}(Z) = Z^{p^{v-2}} = C_{p^2} \times \cdots \times C_{p^2} \times C_p \times \cdots \times C_p,$$

for $i = p^{v-2} + 1, \dots, p^{v-1}$ we get:

$$K_{i,p}(Z) = Z^{p^{v-1}} = C_p \times \cdots \times C_p \quad (\text{just } l \text{ factors}),$$

and, finally, for $i = p^{v-1} + 1$ the series terminates on $K_{i,p}(Z) = \{1\}$.

Therefore, $d = p^{v-1}$ and all the parameters $e(i)$ are zero except the following ones:

$$e(1) = e(p) = \cdots = e(p^{u-2}) = t \quad \text{and} \quad e(p^{u-1}) = l.$$

Thus:

$$\begin{aligned} a &= 1 + (p-1)(t + pt + \cdots + p^{v-2}t + p^{v-1}l) \\ &= 1 + t(p-1) \left(\frac{1-p^{v-1}}{1-p} + l/t \cdot p^{v-1} \right) \end{aligned}$$

and

$$b = (p-1)d = (p-1)p^{v-1}.$$

To deal with the parameters $s(h)$, $h = 1, \dots, c$, for A^β notice that $\gamma_h(A^\beta)$ is a subgroup of the Cartesian power $(\gamma_h(A))^\beta$ (they may not be equal if β is infinite) and, on the other hand, $\gamma_h(A^\beta)$ contains elements with exponent equal to the exponent of $(\gamma_h(A))^\beta$. Therefore, the exponents of $\gamma_h(A^\beta)$ and of $\gamma_h(A)$ are equal for all $h = 1, \dots, c$, and the parameters $s(h)$ are the same for both $\gamma_h(A^\beta)$ and of $\gamma_h(A)$.

By Shield's formula the nilpotency class of $A^\beta \text{Wr } Z$ is the maximum of values

$$(5) \quad ah + (s(h) - 1)b, \quad h = 1, \dots, c.$$

In spite of the fact that having a larger h we get a larger summand ah , it may turn out that for some $h < c$ the exponent of $\gamma_h(A)$ is so much larger than the exponent of

$\gamma_c(A)$ that for the given t the highest value of (5) is achieved not for $h = c$ (examples are easy to build). However, the second summand in (5) may get just c distinct values not dependent on t , whereas the first summand includes a , which grows infinitely and monotonically with t . Thus, even with a fixed l there is a large enough t_0 such that the maximal value of (5) is achieved with $h = c$ for all $t \geq t_0$. To finish this proof just recall that we denoted the exponent of $\gamma_c(A)$ by p^α . $\square$

To introduce our second group we need a finitely generated (and, in fact, also finite, since it is in a locally finite variety) subgroup $\tilde{A}$ of A such that the exponents of terms $\gamma_h(\tilde{A})$ and $\gamma_h(A)$ are equal for each $h = 1, \dots, c$. Clearly, the nilpotency classes of $\tilde{A}$ and of A will then be equal.

Notice that each term $\gamma_h(A)$ contains such an element a_h , the exponent of which is equal to the exponent $p^{s(h)}$ of $\gamma_h(A)$. This is possible, since A is a p -group of finite exponent. Since a_h is an element of the verbal subgroup $\gamma_h(A)$ for the word $\gamma_h(x_1, \dots, x_h)$, there are some finitely many elements $a_{h,1}, \dots, a_{h,r_h} \in A$ such that $a_h \in \gamma_h(\langle a_{h,1}, \dots, a_{h,r_h} \rangle)$. Collecting these finitely many generators for all h , we get the group

$$\tilde{A} = \langle a_{h,i}, \dots, a_{h,r_h} \mid h = 1, \dots, c \rangle,$$

which does have the property we needed. Assume $\tilde{A}$ is a z -generator group and denote by $Y(z, t)$ the product:

$$(6) \quad Y(z, t) = \underbrace{C_{p^v} \times \dots \times C_{p^v}}_{t-z}.$$

Then we have the following value for the nilpotency class of the wreath product $\tilde{A} \text{Wr} Y(z, t)$:

LEMMA 2. *Assume A , $\tilde{A}$, z and α are defined as above. Then there is a positive integer t_1 such that for all $t > t_1$ the nilpotency class of the wreath product $\tilde{A} \text{Wr} Y(z, t)$ is equal to*

$$(7) \quad c + c(t - z)(p - 1) \frac{1 - p^v}{1 - p} + (\alpha - 1)(p - 1)p^{v-1}.$$

PROOF. Denote $Y = Y(z, t)$ and notice that $\tilde{A} \text{Wr} Y$ is nilpotent. Let us compute the K_p -series for Y by the same routine procedure as in previous proof. For $i = 1$ we have $K_{1,p}(Y) = Y$. For $i = 2, \dots, p$ we have:

$$K_{i,p}(Y) = Y^p = C_{p^{v-1}} \times \dots \times C_{p^{v-1}}.$$

For $i = p^k + 1, \dots, p^{k+1}$ we have:

$$K_{i,p}(Y) = Y^{p^{k+1}} = C_{p^{v-(k+1)}} \times \dots \times C_{p^{v-(k+1)}}.$$

In particular, for $i = p^{v-3} + 1, \dots, p^{v-2}$ we get:

$$K_{i,p}(Y) = Y^{p^{v-2}} = C_{p^2} \times \dots \times C_{p^2},$$

for $i = p^{v-2} + 1, \dots, p^{v-1}$ we get:

$$K_{i,p}(Y) = Y^{p^{v-1}} = C_p \times \dots \times C_p,$$

and, finally, for $i = p^{v-1} + 1$ we get $K_{i,p}(Y) = \{1\}$.

Again, $d = p^{v-1}$ and the only non-zero parameters $e(i)$ are:

$$e(1) = e(p) = \dots = e(p^{v-1}) = t - z.$$

Thus:

$$\begin{aligned} a &= 1 + (p-1) \left((t-z) + p(t-z) + \cdots + p^{v-1}(t-z) \right) \\ &= 1 + (t-z)(p-1) \frac{1-p^v}{1-p} \end{aligned}$$

and

$$b = (p-1)d = (p-1)p^{v-1}.$$

Here the parameters $s(h)$, $h = 1, \dots, c$, are the same for $\tilde{A}$ and A , so the nilpotency class of $\tilde{A} \text{ Wr } Y$ is, like in previous proof, the maximum of values

$$(8) \quad ah + (s(h) - 1)b, \quad h = 1, \dots, c,$$

where $p^{s(h)}$ is the exponent of $\gamma_h(A)$. This exponent for some $h < c$ may be so much larger than the exponent of $\gamma_c(A)$ that for the given t the highest value of (8) is achieved not for $h = c$. However, there is a large enough t_1 such that the maximal value of (8) is achieved with $h = c$ for all $t \geq t_1$. Thus, we can assume $h = c$ in formula (8) with $s(c) = \alpha$. $\square$

REMARK 1. Clearly, it would be possible to compute the exact values for the limits t_0 and t_1 in Lemma 1 and Lemma 2. However, that would bring nothing but lengthy calculations, since the exact values of t_0 and t_1 are not relevant for the rest.

Before we proceed to the proof of Theorem 1 let us bring two technical lemmas, where we group a few facts, which either are known in the literature, or are proved by us earlier (see Proposition 22.11 and Proposition 22.13 in [23], Lemma 1.1 and Lemma 1.2 in [16] and also [33]). The proofs can be found in [16], and we bring these lemmas here without arguments:

LEMMA 3. *For arbitrary classes $\mathfrak{X}$ and $\mathfrak{Y}$ of groups and for arbitrary groups X^* and Y , where either $X^* \in \mathbf{Q}\mathfrak{X}$, or $X^* \in \mathbf{S}\mathfrak{X}$, or $X^* \in \mathbf{C}\mathfrak{X}$, and where $Y \in \mathfrak{Y}$, the group $X^* \text{ Wr } Y$ belongs to the variety $\text{var}(\mathfrak{X} \text{ Wr } \mathfrak{Y})$.*

LEMMA 4. *For arbitrary classes $\mathfrak{X}$ and $\mathfrak{Y}$ of groups and for arbitrary groups X and Y^* , where $X \in \mathfrak{X}$ and where $Y^* \in \mathbf{S}\mathfrak{Y}$, the group $X \text{ Wr } Y^*$ belongs to the variety $\text{var}(\mathfrak{X} \text{ Wr } \mathfrak{Y})$. Moreover, if $\mathfrak{X}$ is a class of abelian groups, then for each $Y^* \in \mathbf{Q}\mathfrak{Y}$ the group $X \text{ wr } Y^*$ also belongs to $\text{var}(\mathfrak{X} \text{ Wr } \mathfrak{Y})$.*

Now we can prove the main statement:

PROOF OF THEOREM 1. That the condition of the theorem is sufficient is easy to deduce from the discriminating properties of $C_{p^v}^\infty$ (see [2] or Corollary 17.44 in [23]). Since B and $C_{p^v}^\infty$ generate the same variety $\mathfrak{A}_{p^v}$ and, since $C_{p^v}^\infty$ is isomorphic to a subgroup of B , then by [23, 17.44] B also discriminates $\mathfrak{A}_{p^v}$. It remains to apply Baumslag's theorem: since B discriminates $\text{var}(B) = \mathfrak{A}_{p^v}$, the wreath product $A \text{ Wr } B$ discriminates and, thus, generates the product $\text{var}(A) \mathfrak{A}_{p^v}$ (see [2] or the statements 22.42, 22.43, 22.44 in [23]).

Turning to the proof of necessity of the condition suppose the group B contains no subgroup isomorphic to $C_{p^v}^\infty$. By Prüfer-Kulikov's theorem [30, 10] B is a direct product of some (probably infinitely many) finite cyclic subgroups, the orders of which all are some powers of p . Since B is of exponent p^v , all these orders are bounded by p^v , and there is at least one factor isomorphic to C_{p^v} . By assumption, there are only

finitely many, say l , such factors, and collecting them together, we get $B = B_1 \times B_2$, where

$$B_1 = C_{p^v} \times \cdots \times C_{p^v} \quad (l \text{ factors}),$$

and where B_2 is a direct product of some cycles of orders not higher than p^{v-1} .

Take an arbitrary t -generator group G in variety $\text{var}(A \text{ Wr } B)$. By [23, 16.31] G is in variety generated by all the t -generator subgroups of $A \text{ Wr } B$. Assume T is one of such t -generator subgroups and denote by H its intersection with the base subgroup A^B of $A \text{ Wr } B$. Then

$$T/H \cong (TA^B)/A^B \leq (A \text{ Wr } B)/A^B \cong B$$

and, thus, T is an extension of H by means of an at most t -generator subgroup B' of $B = B_1 \times B_2$. By Kaloujnine-Krasner's theorem [9] the group G is embeddable into $H \text{ Wr } B'$ (see also [25]).

The group B' is a direct product of at most t cycles, of which at most l cycles are of order p^v , and the rest are of strictly lower orders. So B' is isomorphic to a subgroup of $Z(l, t)$ for a suitable t . Since H is a subgroup in A^β , we can apply Lemma 3 and Lemma 4 to get that

$$H \text{ Wr } B \in \text{var}(A^\beta \text{ Wr } Z(l, t)).$$

According to Lemma 1 we get that the nilpotency class of $H \text{ Wr } B$ and of T are bounded by formula (4) for all $t > t_0$.

Our proof will be completed if we discover a t -generator group in $\text{var}(A) \text{ var}(B)$, with nilpotency class higher than (4), at lest for some t .

The group $\tilde{A} \text{ Wr } Y(z, t)$ of Lemma 2 is t -generator, because $\tilde{A}$ is a z -generator group. For sufficiently large $t > t_1$ the nilpotency class of this group is given by formula (7). To compare the values of (4) and (7) notice that they both consist of three summands, of which the first and the third are the same in both formulas. Let us compare the second summands in (4) and in (7). After we eliminate the common multiplier $p - 1$ in both of them, we have:

$$(9) \quad ct \left(\frac{1 - p^{v-1}}{1 - p} + l/t \cdot p^{v-1} \right) = ct \frac{1 - p^{v-1}}{1 - p} + cl p^{v-1}$$

and

$$(10) \quad \begin{aligned} c(t - z) \frac{1 - p^v}{1 - p} &= c(t - z) \left(\frac{1 - p^{v-1}}{1 - p} + p^{v-1} \right) \\ &= ct \frac{1 - p^{v-1}}{1 - p} + ct p^{v-1} - cz \frac{1 - p^{v-1}}{1 - p} - cz p^{v-1}. \end{aligned}$$

The summand $ct \frac{1 - p^{v-1}}{1 - p}$ is the same in (9) and (10), so we can eliminate it also, and just compare the remaining expressions:

$$(11) \quad cl p^{v-1} \quad \text{and} \quad ct p^{v-1} - cz \frac{1 - p^{v-1}}{1 - p} - cz p^{v-1}.$$

Since c, l and v are fixed, the left-hand side of (11) is a positive constant. Since z also is fixed, the second and third summands on the right-hand side of (11) are some negative constants, which make the sum smaller. But, whatever these negative constants be, the other summand $ct p^{v-1}$ in (11) grows infinitely as t grows.

So for sufficiently large t^* the value of (10) is larger than that of (9) (if necessary, we may also take $t^* > t_0, t_1$). Thus, the nilpotency class of the t -generator group

$\tilde{A} \text{ Wr } Y(z, t)$ from the variety $\text{var}(A) \text{ var}(B)$ is higher than the maximum of the nilpotency classes of the t -generator groups in $\text{var}(A \text{ Wr } B)$ for all $t > t^*$. So $\tilde{A} \text{ Wr } Y(z, t)$ does not belong to $\text{var}(A \text{ Wr } B)$, and the proof of the theorem is completed. $\square$

It would not be hard to compute the exact value for t^* in the proof above. We omit it to avoid routine calculations.

REMARK 2. The reader may compare the proofs in this note with proofs in Section 4 in [16] or Section 6 in [18], where we considered similar problem for wreath products of *abelian* p -groups. That time we used the specially defined functions $\lambda(A, B, t)$, and for bounds on nilpotency classes of wreath products of abelian groups we applied Liebeck's formula [12]. As one may notice, in [16, 18] we had a by far simpler situation than what we discussed in Lemma 1 and Lemma 2.

Turning to the examples of usage of Theorem 1 notice that Example 4.6 in [16] and Example 6.9 in [18] already are illustrations of Theorem 1, since they consider wreath products of abelian p -groups of finite exponents. As an example with a nilpotent, non-abelian passive group we may consider:

EXAMPLE 3. The dihedral group D_4 is of nilpotency class 2. Its order is 8 and the exponent is $4 = 2^2$. L.G. Kovács in [11] has computed the variety it generates: $\text{var}(D_4) = \mathfrak{A}_2^2 \cap \mathfrak{N}_2$. That for any finite 2-group B the wreath product $D_4 \text{ Wr } B$ does not generate the product $\text{var}(D_4) \text{ var}(B) = (\mathfrak{A}_2^2 \cap \mathfrak{N}_2) \text{ var}(B)$ is clear from the fact that $D_4 \text{ wr } B$ is a nilpotent group, whereas no product variety may be nilpotent (if both factors are non-trivial). Now take B to be an infinite abelian group of exponent, say, 2^v . By Theorem 1

$$D_4 \text{ Wr } B = (\mathfrak{A}_2^2 \cap \mathfrak{N}_2) \text{ var}(B) = (\mathfrak{A}_2^2 \cap \mathfrak{N}_2) \mathfrak{A}_{p^v}$$

holds if and only if B contains a subgroup isomorphic to $C_{p^v}^\infty$. In particular, if

$$B = \underbrace{C_{p^v} \times \cdots \times C_{p^v}}_n \times \underbrace{C_{p^{v-1}} \times \cdots \times C_{p^{v-1}}}_{\infty} \times \cdots,$$

then $D_4 \text{ Wr } B$ does not generate $(\mathfrak{A}_2^2 \cap \mathfrak{N}_2) \mathfrak{A}_{p^v}$. Moreover, it will not generate it even if we add to B the “large” direct factor

$$\underbrace{C_{p^{v-2}} \times \cdots \times C_{p^{v-2}}}_{\infty} \times \cdots \times \underbrace{C_p \times \cdots \times C_p}_{\infty} \times \cdots.$$

The quaternion group Q_8 of order eight generates the same variety as D_4 (see [23]), and it also is nilpotent of class 2. So a similar example can be constructed for this group also.

References

- [1] G. Baumslag, *Wreath products and p -groups*, Proc. Camb. Philos. Soc. 55 (1959), 224–231.
- [2] G. Baumslag, B.H. Neumann, Hanna Neumann, P.M. Neumann *On varieties generated by finitely generated group*, Math. Z., 86 (1964), 93–122.
- [3] G. Birkhoff, *On the structure of abstract algebras*, Proc. Cambridge Phil. Soc., 31 (1935), 433–454.
- [4] R.G. Burns, *Verbal wreath products and certain product varieties of groups* J. Austral. Math. Soc. 7 (1967), 356–374.

- [5] H. Fitting, *Beitraege zur Theorie der Gruppen endlicher Ordnung* (German), Jahresber. Dtsch. Math.-Ver. 48 (1938), 77–141.
- [6] W. Gaschütz, *Über die Φ -Untergruppe endlicher Gruppen* (German), Math. Z. 58 (1953), 160–170.
- [7] G. Higman, *Some remarks on varieties of groups*, Quart. J. Math. Oxford, (2) 10 (1959), 165–178.
- [8] K.A. Hirsch, *Über lokal-nilpotente Gruppen*, Math. Z. 63 (1955), 290–294 (German).
- [9] L. Kaloujnine, M. Krasner, *Produit complete des groupes de permutations et le problème d’extension des groupes, III*, Acta Sci. Math. Szeged, 14 (1951), 69–82.
- [10] M.I. Kargapolov, Yu.I. Merzlyakov, *Fundamentals of group theory* 4th ed. (Russian), Moscow, Nauka, Fizmatlit 1996. English translation from the 2nd Russian ed. by Robert G. Burns, Graduate Texts in Mathematics 62, New York-Heidelberg-Berlin: Springer-Verlag, XVII 1979.
- [11] L.G. Kovács, *Free groups in a dihedral variety*, Proc. Roy. Irish Acad. Sect. A 89 (1989), no. 1, 115–117.
- [12] H. Liebeck, *Concerning nilpotent wreath products*, Proc. Cambridge Phil. Soc., 58 (1962), 443–451.
- [13] R. Marconi, *On the nilpotency class of wreath products* (Italian), Boll. Unione Mat. Ital., VI. Ser., D, Algebra Geom. 2 (1983), 1, 9–20.
- [14] J.D.P. Meldrum, *On nilpotent wreath products*, Proc. Cambridge Philos. Soc., 68 (1970), 1–15.
- [15] J.D.P. Meldrum, *Wreath products of groups and semigroups*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 74, Harlow, Essex: Longman Group Ltd. xii 1995.
- [16] V.H. Mikaelian, *On varieties of groups generated by wreath products of abelian groups*, Abelian groups, rings and modules (Perth, Australia, 2000), Contemp. Math., 273, Amer. Math. Soc., Providence, RI (2001), 223–238.
- [17] V.H. Mikaelian, *On wreath products of finitely generated abelian groups*, Advances in Group Theory, Proc. Internat. Research Bimester dedicated to the memory of Reinhold Baer, (Napoli, Italy, May-June, 2002), Aracne, Roma, 2003, 13–24.
- [18] V.H. Mikaelian, *Metabelian varieties of groups and wreath products of abelian groups*, Journal of Algebra, 2007 (313), 2, 455–485.
- [19] V.H. Mikaelian, *Varieties generated by wreath products of abelian and nilpotent groups* (Russian and English), Algebra i Logika, 54 (2015), 1, 103–108. Translated in Algebra and Logic, 54 (2015), 1, 70–73.
- [20] V.H. Mikaelian, *The Criterion of Shmel’kin and Varieties Generated by Wreath Products of Finite Groups*, accepted in Algebra i Logika.
- [21] V.H. Mikaelian, *A Classification Theorem for Varieties Generated by Wreath Products of Groups*, in preparation.
- [22] V.H. Mikaelian, A.Yu. Olshanskii *On abelian subgroups of finitely generated metabelian groups*, Journal of Group Theory, 16 (2013), 695–705.
- [23] Hanna Neumann, *Varieties of Groups*, Varieties of groups (Ergebn. Math. Grenz., 37), Berlin-Heidelberg-New York, Springer-Verlag 1967.
- [24] A.Yu. Olshanskii *The Neumanns-Shmel’kin’s theorem (Russian)*, Vestnik Mosk. Univ., Ser. Matem.(1986), N 6, 61–64.
- [25] A.Yu. Olshanskii, *On Kaluzhnin-Krasner’s embedding of groups*, to appear.
- [26] B.I. Plotkin, *On some criteria of locally nilpotent groups*, Uspehi Mat. Nauk 9 (1954), 181–186. Translated in Amer. Math. Soc. Translations (2) 17 (1961), 1–7.
- [27] B.I. Plotkin, *Radical groups*, Mat. Sb. 37 (1955), 507–526. Translated in Amer. Math. Soc. Translations (2) 17 (1961), 9–28.
- [28] B.I. Plotkin, *Generalized soluble and nilpotent groups*, Uspehi Mat. Nauk 13 (1958), 89–172. Translated in Amer. Math. Soc. Translations (2) 17 (1961), 29–115.
- [29] D.J.S. Robinson, *Finiteness Conditions and Generalized Soluble Groups II*, Springer-Verlag, Berlin 1972.

- [30] D.J.S. Robinson, *A Course in the Theory of Groups*, second edition, Springer-Verlag, New York, Berlin, Heidelberg 1996.
- [31] D. Shield, *Power and commutator structure of groups*, Bull. Austral. Math. Soc., 17 (1977), 1–52.
- [32] D. Shield, *The class of a nilpotent wreath product*, Bull. Austral. Math. Soc., 17 (1977), 53–89.
- [33] A.L. Shmel'kin, *Wreath products and varieties of groups*, Izv. AN SSSR, ser. matem., 29 (1965), 149–170 (Russian). Summary in English: Soviet Mathematics. Vol. 5. No. 4 (1964). Translation of Dokl. Akad. Nauk S.S.S.R. for Am. Math. Soc.

Yerevan State University, Alex Manoogian 1, Yerevan 0025, Armenia.

American University of Armenia, 40 Marshal Baghramyan Ave., Yerevan 0019, Armenia.

E-mail addresses: v.mikaelian@gmail.com, vmikaelian@ysu.am, vmikaelian@aua.am.